


Two tiny beetle fossils offer evolution and biogeography clues

January 18 2019



Sphaerotherax uenoi from mid-Cretaceous Burmese amber. Credit: NIGPAS


It is well-known that living fossils exhibit stasis over geologically long time scales. Examples are the panda and ginkgo. Now, two tiny beetles

trapped in 99-million-year-old amber may join this group.

Beetles are the most species-rich group of animals on our planet. Among four extant suborders of beetles, polyphaga is the largest and most diverse group. The origin and early evolutionary history of polyphagan beetles have been largely based on evidence from the derived and diverse core polyphaga, whereas little is known about the species-poor basal polyphagan lineages, which include Clambidae and four other extant families.

Recently, an international team led by Dr. Cai Chenyang, from the Nanjing Institute of Geology and Palaeontology of the Chinese Academy of Sciences, reported two new and rare species of the extant family Clambidae from Burmese amber: *Acalyptomerus thayerae* and *Sphaerotherax uenoi*. They are important for understanding the early evolution and biogeography of the family and even for polyphagan beetles.

Clambidae is a small group of small-sized (usually 0.7-2.0 mm long) polyphagan [beetles](#) distributed worldwide, with approximately 150 described species grouped in five extant genera. Most clambid adults occur in decaying vegetation, [leaf litter](#) and rotten wood and occasionally fly at dusk.



Geographical distribution of *Acalyptomerus thayerae* and *Sphaerotherax uenoi* and their related extant counterparts. Credit: NIGPAS

The two new species, represented by five well-preserved fossils, were discovered by Dr. Cai and colleagues from mid-Cretaceous Burmese amber. Both species are extremely morphologically close to their living counterparts, and can be placed in extant genera. *Acalyptomerus thayerae* (about 1.05-1.15 mm long) has a close affinity to *A. herbertfranzi*, a species currently occurring in Mesoamerica and northern South America. *Sphaerotherax uenoi* (about 0.71 mm long) is closely related to extant species of *Sphaerotherax*, which are usually collected in forests of *Nothofagus* in Australia, Chile and New Zealand.

The discovery of two Cretaceous species from northern Myanmar indicates that both genera had lengthy evolutionary histories, originating at least by the earliest Cenomanian, and were probably more widespread than at present.

Remarkable morphological similarities between fossil and living [species](#) suggest that both genera changed little over long periods of geological time, which is usually considered to be a feature of living fossils. The long-term persistence of similar mesic microhabitats such as leaf litter may account for the 99-million-year morphological stasis in *Acalyptomerus* and *Sphaerotherax*.



Acalyptomerus thayerae from mid-Cretaceous Burmese amber. Credit: NIGPAS

More information: Chenyang Cai et al, Basal polyphagan beetles in mid-Cretaceous amber from Myanmar: biogeographic implications and long-term morphological stasis, *Proceedings of the Royal Society B: Biological Sciences* (2019). [DOI: 10.1098/rspb.2018.2175](https://doi.org/10.1098/rspb.2018.2175)

Provided by Chinese Academy of Sciences

Citation: Two tiny beetle fossils offer evolution and biogeography clues (2019, January 18)
retrieved 27 September 2024 from
<https://phys.org/news/2019-01-tiny-beetle-fossils-evolution-biogeography.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--